



ECOLOGICAL SOCIETY OF AMERICA

Ecology/Ecological Monographs/Ecological Applications

PREPRINT

This preprint is a PDF of a manuscript that has been accepted for publication in an ESA journal. It is the final version that was uploaded and approved by the author(s). While the paper has been through the usual rigorous peer review process of ESA journals, it has not been copy-edited, nor have the graphics and tables been modified for final publication. Also note that the paper may refer to online Appendices and/or Supplements that are not yet available. We have posted this preliminary version of the manuscript online in the interest of making the scientific findings available for distribution and citation as quickly as possible following acceptance. However, readers should be aware that the final, published version will look different from this version and may also have some differences in content.

The doi for this manuscript and the correct format for citing the paper are given at the top of the online (html) abstract.

Once the final published version of this paper is posted online, it will replace the preliminary version at the specified doi.

1 Running head: **Resistance to floods**

2 Title: **Quantifying invertebrate resistance to floods: a global-scale meta-analysis**

3

4

Laura E. McMullen* and David A. Lytle

5

6 *Department of Zoology, Oregon State University, 3029 Cordley Hall, Corvallis, Oregon 97331*

7 *USA*

8

9

10

11 ** Corresponding author. Current address: Laura E. McMullen, ICF INTERNATIONAL, 615 SW*

12 *Alder St., 2nd Floor, Portland, OR 97205*

13 *Email: Laura.McMullen@icfi.com*

14

15 *Email: lytle@science.oregonstate.edu*

16 **Abstract**

17 Floods are a key component of the ecology and management of riverine ecosystems
18 around the globe, but it is not clear whether floods have predictable effects on organisms that can
19 allow us to generalize across regions and continents. To address this, we conducted a global-
20 scale meta-analysis to investigate effects of natural and managed floods on invertebrate
21 resistance, the ability of invertebrates to survive flood events. We considered 994 studies for
22 inclusion in the analysis, and after evaluation based on *a priori* criteria, narrowed our analysis to
23 41 studies spanning 6 of the 7 continents. We used the natural log ratio of invertebrate
24 abundance before and within 10 days after flood events because this measure of effect size can
25 be directly converted to estimates of percent survival. We conducted categorical and continuous
26 analyses that examined the contribution of environmental and study design variables to effect
27 size heterogeneity, and examined differences in effect size among taxonomic groups. We found
28 that invertebrate abundance was lowered by at least half after flood events. While natural vs.
29 managed floods were similar in their effect, effect size differed among habitat and substrate
30 types, with pools, sand, and boulders experiencing the strongest effect. Although sample sizes
31 were not sufficient to examine all taxonomic groups, floods had a significant, negative effect on
32 densities of Coleoptera, Eumalacostraca, Annelida, Ephemeroptera, Diptera, Plecoptera, and
33 Trichoptera. Results from this study provide guidance for river flow regime prescriptions that
34 will be applicable across continents and climate types, as well as baseline expectations for future
35 empirical studies of freshwater disturbance.

36 **Key words**

37 River management, environmental flows, quantitative synthesis, disturbance ecology

38

esa
preprint

39 **Introduction**

40 Freshwater is becoming an increasingly important and scarce resource around the world
41 (Yeston et al 2006). While humans have altered freshwater ecosystems through damming in the
42 majority of large-river systems in the world (Nilsson et al. 2005), there is a trend to bring flows
43 back to a more natural regime and to recognize rivers themselves as legitimate users of water
44 (Naiman et al. 2002). Environmental flows are one paradigm used to manage rivers across the
45 world, with over 200 different methodologies having been developed (Tharme 2003). Under this
46 broad framework, elements of the natural flow regime are mimicked to produce desired
47 ecological outcomes, such as increased biodiversity or habitat creation for target species.

48 Despite the diversity of methods that have been developed at various scales to prescribe
49 environmental flows to rivers (Jowett 1997, Arthington et al. 2006), there is little quantitative
50 information regarding how flood events affect specific biota and ecosystem processes (Bunn and
51 Arthington 2002). This quantitative information is necessary for accurate parameterization of
52 predictive models of ecological effects of managed flow regimes, and can aid in forming useful
53 hypotheses for further scientific studies on freshwater ecology.

54 Overall, while there are many case studies investigating effects of floods on aquatic
55 organisms, differences in river type, regional climate, and continental setting make it difficult to
56 draw general conclusions (Resh et al. 1988, Death 2010). A quantitative understanding of how
57 aquatic organism populations immediately respond to disturbance events would lead to better
58 predictions of post-flood population sizes, simpler interpretation of post-flood monitoring data,
59 and a better understanding of organisms' responses to disturbance events (Poff and Zimmerman
60 2010).

61

62 In this study we used a global-scale meta-analytic study to examine the quantitative
63 relationships between flood events and change in invertebrate abundance (resistance). We
64 focused on aquatic invertebrates because they encompass a wide array of life-history and
65 behavioral characteristics that can inform studies of other aquatic taxa. Specifically, our goals
66 were to 1) determine whether effects of natural versus prescribed flood events differ and to what
67 degree, 2) investigate differences in effects of floods among riverine habitat types and study
68 designs, 3) determine whether a flood's relative magnitude affects organism resistance, and 4)
69 explore differences in response to flooding across taxonomic groups.

70 **Methods**

71 *Literature search*

72 We searched the literature with *a priori* criteria for appropriate primary case studies
73 concerning effects of floods on aquatic invertebrate abundance immediately after flood events.
74 We used the electronic database Web of Science (including papers from 1970-2010) to identify
75 potential studies for inclusion. We used the terms *spate* or *flood*, *macroinvertebrate* or *macro-*
76 *invertebrate* or *insect* or *invertebrate*, and *benthic* or *aquatic* or *stream* as keywords, resulting in
77 994 potential studies. We evaluated each study for inclusion with the following criteria. Studies
78 were required to be primary research papers, and needed to contain information on independent
79 flood events in rivers, streams, or artificial stream channels, with both pre- and post- data on
80 aquatic invertebrate density in relation to floods (e.g., invertebrate abundance per square meter,
81 or abundance per cage, artificial substrate, or rock). We excluded studies that only reported
82 correlation coefficients or significance values concerning flood effects on invertebrates. We also
83 excluded studies that had confounding treatments such as insecticide application. We included
84 both natural and managed floods. The pre-flood samples must have occurred within 60 days of

85 the flood event, and the post-flood samples within 10 days of the flood event. If other papers
86 were cited that could contain needed, missing information, we included data from those papers as
87 well. With these criteria in place, we obtained 41 studies for analysis (Table 1).

88 We collated data from these studies in two ways, each intended to test different questions
89 about invertebrate response to flood events (Table 2):

90 1) General data set. Total abundance of all invertebrates per unit area, without respect
91 to taxonomy, was used as the sample unit. This conservative approach avoids the
92 issue of independence among taxa at a given site, but fails to identify taxon-specific
93 differences in flooding response.

94 2) Taxon-specific data set. Abundance of different taxonomic groups of invertebrates
95 per unit area, broken down by lowest taxonomic level reported in studies, represents
96 the sample unit. Within a study, taxonomic groups were weighted equally. This
97 approach allowed us to identify potential taxon-specific differences in flooding
98 response.

99 For example, a study could have reported abundance before and after a flood event for
100 five taxa. For the general data set, we would sum the abundances of the five taxa and consider
101 this a sample unit. For the taxon-specific data set, the abundance before and after the flood event
102 for each of the five taxa was considered a sample unit. In this scenario, we would have obtained
103 one sample unit for the general data set, and five sample units for the taxon-specific data set.

104 These alternative replication schemes have different implications for the interpretation of results.

105 For the general data set, the cumulative effect size (Rosenberg et al. 2000) of floods on
106 total invertebrate abundance could be biased towards taxa that generally occur in higher
107 abundance. For the taxon-specific data set, the cumulative effect size is representative of the

108 overall magnitude of the effect of floods on all taxa treated as individual units of replication in
109 all the studies in the data set. Besides calculating a cumulative effect size of floods on overall
110 invertebrate abundance from the taxon-specific data set (and using this value in categorical and
111 continuous analyses), we were also able to compare effect of floods among different taxonomic
112 groups.

113 For the general data set, if a study reported the total invertebrate densities before and after
114 the flood event, these numbers were used. If a study only reported densities for specific taxa,
115 densities of individual taxa were aggregated so long as data for three or more orders of
116 invertebrates were reported (Table 2). For the taxon-specific data set, we first recorded
117 invertebrate data at the finest taxonomic level reported in each study, and then standardized to
118 higher taxonomic levels where appropriate. We considered different taxonomic groups within a
119 study independently. For taxon-specific analyses, we also included studies in which data were
120 reported as a percent change from pre to post-flood.

121 Within the taxon-specific data set, data were standardized to different taxonomic levels
122 depending on the analysis being performed. For analyses that were performed using both the
123 general data set and the taxon-specific data set, sample units consisted of abundances for each
124 insect Order (and other levels for non-insects). Thus, data were standardized to this level by
125 summation of lower taxonomic levels (if the data were reported as density data) or by averaging
126 (if the data were reported as a percent change). A categorical analysis among groups of taxa at
127 these higher-level taxonomic groupings was also performed.

128 A second set of taxon-specific analyses were conducted at the family level. All groups of
129 taxa determined in the first set of taxon-specific analyses were analyzed for inclusion in this next
130 step of analysis. For a group of invertebrates to be included, it had to have sub-group data for at

131 least 2 disparate groups at the next classification level with $n \geq 5$ for each, and with data derived
132 from at least 3 separate studies for each sub-group. The goal of this set of analyses was to
133 determine whether significant differences in resistance to flooding can be detected among groups
134 at finer classification levels.

135 We included data only for flood events at least 60 days apart, with no significant floods
136 within 60 days prior to the flood event, for each river in each study. We included data for
137 multiple sites per river per study, if data were reported for multiple longitudinal sites. Although
138 including multiple flood events and longitudinal river sites from a single study in the analysis
139 could cause a lack of spatial or temporal independence, this is a common problem in meta-
140 analysis, and we concluded that exclusion of these data would be too great of an information
141 loss. If data from multiple rivers were reported in a study, we included data from all rivers in the
142 analyses. When needed, we used Data Thief III software (Tummers 2006) to extract data from
143 graphs.

144 *Examining resistance via effect size*

145 Resistance can be defined as the ability of a population or community to withstand a
146 disturbance event (*sensu* Grimm and Fisher 1989) so we calculated effect size of floods on
147 aquatic invertebrate taxa within 10 days after the flood event. The primary response variable of
148 interest was density of invertebrate taxa per unit area. We used natural log response ratio (R) as
149 the measure of effect size in this study: $\ln(\text{density of invertebrates post-flood} / \text{density of}$
150 $\text{invertebrates pre-flood})$. Thus, a negative effect size indicated a reduction in density of
151 individuals following a flood event. Taking the natural log of the response ratio linearizes the
152 results by equally accounting for the numerator and denominator, and normalizes the sampling
153 distribution of the response ratio (Hedges et al. 1999).

154 *Meta-analytic techniques*

155 We performed an unweighted analysis, as 7 studies did not report variance and would
156 have been excluded from the analysis. Additionally, summation of invertebrate data from lower
157 to higher taxonomic levels for standardization disallowed accounting for variance. We used an
158 unstructured and unweighted random effects model in MetaWin (Rosenberg et al. 2000) to
159 evaluate overall effect size of floods on aquatic invertebrates. Effect sizes, in the case of ln
160 response ratio, are considered significant if their 95% confidence intervals do not overlap zero
161 (Rosenberg et al. 2000; Shurin et al. 2002).

162 Using both the general data set and the highest-aggregated level of the taxon-specific data
163 set, we examined resistance of overall invertebrate density to flood events, and also explored
164 potential effects of natural versus managed floods, habitat type, substrate type, collection
165 method, and whether the flood happened in a month with higher or lower average rainfall with
166 categorical analyses. We also performed an analysis of resistance of invertebrates as a function
167 of the number of days since the flood event, and as a function of the relative flood magnitude
168 (peak discharge/ mean discharge or mean baseflow). Continuous analyses were performed as
169 unweighted linear regressions.

170 We reported all statistics at the $\alpha=0.05$ significance level. We performed the majority of
171 analyses using MetaWin (Rosenberg et al. 2000), and we also used SigmaPlot (SigmaPlot 2004)
172 for data visualization and some analyses. For categorical analyses, we included categories only
173 if the number of sample units in a given category ≥ 5 , and if the sample units were derived from
174 at least 3 separate studies. When we detected a significant difference between categories,
175 unplanned comparisons of means were conducted using the Tukey-Kramer method (Sokal and
176 Rohlf 2000).

177 We examined a funnel plot of effect size vs. sample size to detect publication bias, such
178 as underreporting of non-significant studies. Assuming no publication bias, smaller sample sizes
179 are expected to have greater error spread, the cumulative effect size is expected to be
180 independent of sample size, and normal distribution of individual studies is expected at all
181 sample sizes (Palmer 1999).

182 **Results**

183 The 41 studies included in the analyses spanned 13 countries and 37 rivers, streams, or
184 stream systems (Table 1). There appeared to be slight asymmetry in the funnel plots of both the
185 general and taxon-specific data sets, indicating that there could be a relationship between
186 treatment effect and sample size, but there is not enough evidence to indicate strong publication
187 bias. Smaller samples sizes had greater error spread as expected. Especially for the taxon-
188 specific data set, distribution of effect sizes seemed to have a longer left (negative) than right tail.
189 This could be because floods generally have a negative effect on invertebrate abundance, and
190 thus the left tail of the distribution was more prominent. However, it could be due to some
191 under-reporting of studies where floods had positive effects on invertebrate abundance, and these
192 different potential underlying reasons cannot be teased apart.

193 *Overall effect*

194 Using the general data set, there was a significant, negative effect of floods on the overall
195 density of invertebrates within 10 days of a flood event (cumulative effect size -1.01, 95% CI (-
196 1.27 to -0.76), $n=90$) (Figure 1). This is equivalent to a reduction of 53-72% of overall density
197 of invertebrates within 10 days of a flood event. To check for independence, we ran the same
198 analysis on a data set with one sample unit randomly selected from each study and found a

199 significant, negative effect that is not significantly different from the effect calculated from the
200 full data set (cumulative effect size -0.8506, 95% CI (-1.1074 to -0.5938), $n=34$).

201 For the taxon specific data set, there was also a significant, negative effect of floods on
202 the overall density of invertebrates within 10 days of a flood event (cumulative effect size -1.15,
203 95% CI (-1.37 to -0.93), $n=340$). This is equivalent to a reduction of 61 to 75% of individuals in
204 all groups of invertebrates within 10 days of a flood event.

205 *Categorical analyses*

206 Using the general data set, effect size of floods on invertebrate density did significantly
207 differ between habitat types ($P<0.01$, groups=3, Figure 1, Table 3). Invertebrates were most
208 severely reduced by floods in pool habitats, which differed significantly in effect size from run or
209 riffle habitats (Table 3). Using the taxon specific data set, invertebrates were again most
210 severely reduced by floods in pool habitats, while they were least reduced in run habitats
211 ($P=0.003$, groups=3, Figure 1, Table 3), and in this case all three habitats had significantly
212 different effect sizes from each other (Table 3).

213 There was no significant difference found between effect size of natural versus managed
214 floods on invertebrate density using the general data set ($P=0.98$, groups=2) or the taxon
215 specific data set ($P=0.4$, groups=2, Figure 1, Table 3). There also was no significant difference
216 in effect size between collection methods using the general data set ($P=0.12$, groups=5, Table 3)
217 or the taxon specific data set ($P=0.17$, groups=5, Figure 1, Table 3).

218 Using the general data set no significant difference in effect size between invertebrate
219 densities collected from different substrate types was detected ($P=0.63$, groups=6, Figure 1,
220 Table 3). However, using the taxon specific data set, complex differences in effect size among
221 substrate types were found ($P=0.003$, groups=6, Figure 1, Table 3), with invertebrate density

222 being most reduced in sandy substrates and least reduced on wood. There was also no
223 significant difference in effect size between floods that happened in a typical ‘wet’ month
224 (higher than mean annual rainfall) or ‘dry’ month (lower than mean annual rainfall) using the
225 general data set ($P=0.51$, groups=2, Table 3) or the taxon specific data set ($P=0.68$, groups=2,
226 Figure 1, Table 3).

227 *Continuous analyses*

228 A continuous model analysis showed that effect size became smaller in magnitude (closer
229 to zero) with days since flood event (slope $P=0.02$, $n=89$) (Figure 2). However, with removal of
230 the outlier with the largest effect size at 10 days post-flood, the relationship was no longer
231 significant (slope $P=0.11$, $n=88$). A continuous model analysis using the taxon-specific data set
232 showed no significant effect of days since flood on effect size within 10 days of a flood event
233 (slope $P=0.9$, $n=339$).

234 When including all data from all river and habitat types in a continuous model analysis of
235 effect size versus relative flood magnitude, there was no significant trend detected. However,
236 when a continuous model analysis was performed using only samples from riffle or run habitats
237 composed of primarily cobble or gravel substrate (generalized habitat types that were most
238 commonly reported on in primary studies), effect size became greater with increasing relative
239 flood magnitude (slope $P<0.01$, $n=49$) (Figure 3). As with the general data set, when including
240 all data there was no significant effect of relative flood magnitude on effect size. There was a
241 significant increase in effect size with relative flood magnitude when examining only riffle or
242 run habitats dominated by cobble or gravel substrate (slope $P<0.0001$, $n=202$). It is possible
243 that there is a threshold at a relative flood magnitude of approximately 40-50, where the response
244 to flooding is suddenly much stronger.

245 *Taxon-specific analyses of resistance*

246 Floods had a significant, negative effect on densities of Coleoptera, Eumalacostraca,
247 Annelida, Ephemeroptera, Diptera, Plecoptera, and Trichoptera (95% confidence intervals did
248 not overlap zero, Figure 4). Floods did not have a significant effect on densities of Acari,
249 Mollusca, or Platyhelminthes (95% confidence intervals did overlap zero, Figure 4). However,
250 there were no significant categorical differences between groups, since all of their confidence
251 intervals overlapped ($P=0.26$, Table 3).

252 Application of selection criteria for categorical analyses at finer taxonomic levels
253 narrowed the groups for further analysis to Diptera, Ephemeroptera, Plecoptera, and Trichoptera.
254 Of these groups, categorical analyses only found significant differences among families within
255 each order for the Diptera, with Chironomidae experiencing significantly greater post-flood
256 reduction than Tipulidae or Simuliidae ($P=0.049$, $n=4$, Figure 4, Table 3). All mayfly families
257 experienced significant reduction following flood events.

258 **Discussion**

259 This meta-analysis found a significant reduction in overall invertebrate abundance and a
260 reduction in abundance of major groups of invertebrates immediately after flood events in rivers.
261 This relationship was apparent despite large differences in river type (parent geology, gradient,
262 catchment size), regional climate, and continental setting. While a number of case studies exist
263 concerning prescribed high flow releases and ecosystem effects, and other papers have published
264 information on natural floods and effects on invertebrates, there is a paucity of among-stream
265 studies of flood effects on aquatic invertebrates (Death 2007). This is the first calculation of
266 values for immediate invertebrate reduction after floods across studies at a global scale.

267 There is a need for increased ability to predict outcomes of river flow management on
 268 aquatic biota (Death 2007, Souchon et al. 2008, Poff 2009). While some studies have considered
 269 quantitative, cross-system effects of river flow management on aquatic organisms and
 270 communities (Bickford and Skalski 2000, Monk et al. 2006, Haxton and Findlay 2008, Stewart et
 271 al. 2009), this study contributes new information to our growing synthetic knowledge.

272 One purpose of meta-analyses is to generate predictive hypotheses for further
 273 experimentation and evaluation (Osenberg et al. 1999, Lajeunesse 2010). Because log response-
 274 ratios may be easily translated into percent reductions, the overall effect size of density change of
 275 invertebrates due to floods, and other quantitative data regarding effect sizes in this study, may
 276 be used directly for modeling or quantitative prediction of management outcomes. The results of
 277 this meta-analysis can therefore be used to predict responses of biota to flood events and to
 278 parameterize general models of flood effects on aquatic organism abundance.

279 *What is the overall estimate of reduction of invertebrates post-floods, and does this differ among*
 280 *natural versus managed floods?*

281 The overall values of resistance from both data sets are in concordance and show that
 282 invertebrates are generally reduced in numbers by at least half immediately after flood events,
 283 and we found no evidence for differing effects of natural versus managed floods on invertebrate
 284 resistance. While lack of evidence for a statistical relationship does not necessarily mean that a
 285 relationship does not exist, our results indicate that as far as we know, general inferences drawn
 286 from mensurative (natural) flood experiments may be applied to development of manipulative
 287 flood experiments (Konrad et al. 2012). While mensurative flow experiments do not have true
 288 replication, pre-condition standardization, or control of treatment size (Konrad et al. 2012), they
 289 are useful in the context of synthesis of data from multiple, observable, quantified studies.

290 However, managed floods can sometimes differ from natural floods in ways that can
291 affect the response of organisms. For example, some aquatic invertebrates use proximate cues
292 such as rainfall or flow to escape from floods or return to the stream post-flood (Lytle et al. 2008,
293 Lytle & White 2007). If a managed flood lacks these proximate cues, or follows a hydrograph
294 pattern that is not typical of natural floods (e.g., abrupt increases or decreases in flow), the
295 organisms could be negatively affected.

296 *How do environmental variables influence heterogeneity in effect of floods on invertebrate*
297 *resistance?*

298 Categorical analysis of both data sets demonstrated significant differences in effect of
299 floods on invertebrate resistance among different general habitat types. While one data set
300 showed differences among all three habitat types- riffle, run, and pool- the other showed that
301 only pool habitats differed from riffle and run habitats. In general, pool invertebrates were
302 reduced in density to a greater degree than invertebrates in riffles or runs. There is evidence that
303 substrates in pools are more easily scoured by spates than substrates in riffles or runs
304 (Scarsbrook and Townsend 1993, LaPointe et al. 2000, Harrison and Keller 2007). This could
305 also affect the egg or larval stages of other aquatic organisms, such as salmon redds. Eggs in
306 riffles or run likely have a higher chance of withstanding high flow events than those in pool
307 habitats. Aquatic macrophytes in riffle or run habitats may also be less susceptible to flow events.
308 These are hypotheses worth testing further.

309 Substrate type was a significant factor when categorically examining differences in effect
310 sizes from the taxon-specific data set, but not when using the general data set. Differences
311 among groups demonstrated by the taxon-specific data set were complex, with invertebrates
312 reduced to the greatest degree on boulder and sand substrates, and least reduced on wood

313 substrates. Wood and cobble can act as a refuge for invertebrates during flood events by
314 providing greater structural complexity (Hax and Golladay 1998, Palmer et al. 1996). Sand, the
315 smallest-diameter substrate evaluated here, would be moved by the least force and thus be the
316 most easily disturbed of these substrates. Boulders, one of the larger substrates analyzed, also
317 showed very low resistance of invertebrates. This may be due to the lack of interstitial spaces on
318 boulders to act as refuges (Lancaster 1992), or the frequent covering of boulders with silt and
319 associated algae or macrophytes which may be easily disturbed by floods. Intermediate-sized
320 substrates may provide the most protection for invertebrates from flood events. These results are
321 also important for egg and larval stages of other aquatic organisms (fish, amphibians) and small
322 adult fish or amphibians, which may also withstand flood events best on intermediate substrate.
323 The specific habitat sampled, its constituent substrate, and how it was sampled must be taken
324 into account when predicting flood effects on organisms, due to the great differences in
325 resistance these variables confer on the organisms.

326 *Is there evidence for 'hidden resistance,' or a short-term increase in invertebrate abundance*
327 *post-flood?*

328 Analysis of the general data set showed that invertebrates significantly increased in
329 numbers within 10 days after a flood event, although with removal of an extreme data point this
330 relationship was no longer significant. Although succession via recolonization and recruitment
331 may begin immediately after flooding, the evident increase in resistance of invertebrates within
332 10 days of a flood event may encompass 'hidden survival' since the majority of stream-dwelling
333 organisms have life-cycles greater than 10 days. Organisms may be displaced by the flood into
334 marginal habitats (side channels, deep pools) or buried by substrates. Indeed, invertebrates in
335 several groups have the ability to return to the active stream channel if displaced by a flood

336 (Lytle et al. 2008), and still other taxa are known to abandon streams prior to flooding and
337 eventually return (Lytle & White 2007, Lytle 2000). Thus, we cannot assume that low incidence
338 of organisms directly after flood events is always indicative of mortality. Examining short-term
339 recovery of longer-lived aquatic organisms, including fish and amphibians, directly after flood
340 events might provide more evidence for ‘hidden survival’. This has important implications for
341 monitoring events after floods, as monitoring too quickly after a flood event could over-estimate
342 mortality.

343 Analysis of the taxon-specific data set showed no relationship between effect size and
344 days since event in a continuous model analysis. With such varied life-history patterns and
345 overall lifespans in aquatic invertebrates, what is defined as ‘resistance’ versus ‘resilience’ may
346 vary between groups. For example, fast life-cycled mayflies such as *Fallceon quilleri*
347 (Ephemeroptera: Baetidae) may transform from egg to reproductive aerial adult in as fast as 7
348 days (Gray 1981), and their aerial stage can escape river-bed flood events. Measuring resistance
349 of this species to floods may need to happen within a day or two of a flood event, as their
350 populations may immediately rebound immediately after flood events. For longer-lived
351 organisms, and those without aerial stages, the effects of flood disturbance may be evident for a
352 much longer time period.

353 *How does flood magnitude influence invertebrate resistance?*

354 When including all data, both for the general data set and the taxon-specific data set,
355 there were no significant changes in effect sizes with relative flood magnitude. However, for
356 some specific habitats (riffles, runs; cobble or gravel substrates) we did find an effect. We
357 believe that flood magnitude does play an important role in shaping the effect of floods on
358 invertebrates and other aquatic organisms, and that the effect of flood magnitude on invertebrates

359 was masked in our full data set because it spanned such a wide array of habitats that differed in
360 response to flooding. Thus, any broad generalizations about the effect of floods on invertebrates
361 must still account for differences in response due to habitat and substrate type.

362 *Does resistance to floods differ among taxonomic groups?*

363 While there was no significant categorical difference between groups at the level of
364 Order (insects) and higher (non-insects), some groups were significantly affected by flood events
365 (95% confidence intervals not overlapping zero), while others were not (95% confidence
366 intervals overlapping zero). All insect groups were significantly affected by flood events. The
367 only groups not shown to be significantly affected were water mites (Acari), molluscs
368 (Mollusca), and flatworms (Platyhelminthes). However, variance in effect size within these
369 groups was also very large, and sample sizes were low, so this may be an issue of statistical
370 power rather than biological response. Similar analyses could potentially be performed by trait
371 group instead of by taxonomic categories, which could answer questions about which
372 morphological, life-history, or behavioral traits are most successful at providing organisms
373 defense against flood disturbance events. However, information on lower levels of taxonomic
374 organization for reported invertebrates would likely be needed since traits may vary widely at
375 higher taxonomic levels.

376 There were not enough data reported on some aquatic insect taxa (and other aquatic
377 invertebrates) to justify including them. These less-commonly reported insect groups included
378 odonates (dragonflies and damselflies), hemipterans (true bugs), megalopterans (alderflies and
379 dobsonflies), collembolans (springtails), and aquatic lepidopterans (moths). Many studies
380 reported only a subset of taxa, generally those found to be most abundant in the system. Greater
381 reporting of data regarding all taxa collected and identified instead of just the most abundant taxa

382 collected would broaden our ability to discern the generalities critical to both basic biological
383 understanding and effective management. Also, there were few available published studies from
384 1970-2010 quantifying immediate effects of floods on biota from Africa, Asia, Central and South
385 America. In fact, all together only 13% of rivers and streams reported on in this analysis are
386 drawn from those continents, while 49% were in the United States and Canada. More studies
387 concerning flows in these under-reported countries are needed.

388 This meta-analysis suggests further studies which would be useful to answer specific
389 questions concerning disturbance effects on aquatic organisms. For example, organisms
390 inhabiting pool versus riffle or run habitats in rivers could be censused to determine if
391 differences in community structure exist. If so, it could be examined whether these organisms
392 inherently differed in ability to survive floods, regardless of initial habitat preference, or whether
393 organisms in pools are simply more susceptible due to greater scouring. This could be useful in
394 predicting outcomes of direct management of riverine morphology on aquatic populations, i.e.
395 influences of artificial enhancement of pools via additions of boulders or wood. Streamside
396 experiments could be undertaken to closely examine the influence of substrate type on flood
397 effects. Populations of specific taxa could be closely tracked after flood events to elucidate
398 whether resistance measurements may be influenced by short-term 'hidden resistance'. Also,
399 comprehensive, quantitative evaluation of other aspects of the flow regime (drought, base flows,
400 timing of flow events, etc.) and studies on other organisms would be useful to solidifying a
401 scientific framework on which to base specific prescribed flow events and to predict ecological
402 reactions to climate induced hydrologic changes.

403

404 **Acknowledgements**

405 Funding was provided by a NSF Predoctoral Fellowship to L.E.M. Helpful comments
406 that resulted in substantial improvement to this manuscript were provided by Kate Boersma,
407 Elizabeth Borer, and several anonymous reviewers.

esa
preprint

408 **Literature cited**

409 Angradi, T.R. 1997. Hydrologic context and macroinvertebrate community response
 410 to floods in an Appalachian headwater stream. *American Midland Naturalist*
 411 138: 371-386.

412 Arthington, A. H., S. E. Bunn, N. L. Poff, and R. J. Naiman. 2006. The challenge of
 413 providing environmental flow rules to sustain river ecosystems. *Ecological*
 414 *Applications* 16: 1311-1318.

415 Baumgartner, A., and J.A. Waringer. 1997. Longitudinal zonation and life cycles of
 416 macrozoobenthos in the Mauerbach near Vienna, Austria. *Internationale*
 417 *Revue gesamten Hydrobiologie* 82: 379-394.

418 Bickford, S. E., and J. R. Skalski. 2000. Reanalysis and interpretation of 25 years of
 419 Snake-Columbia river juvenile salmonid survival studies. *North American*
 420 *Journal of Fisheries Management* 20: 53-68.

421 Bond, N. R., and B.J. Downes. 2000. Flow-related disturbance in streams: an
 422 experimental test of the role of rock movement in reducing macroinvertebrate
 423 population densities. *Marine and Freshwater Research* 51: 333-337.

424 Boulton, A. J., C. G. Peterson, N. B. Grimm, and S. G. Fisher. 1992. Stability of an
 425 aquatic macroinvertebrate community in a multiyear hydrologic disturbance
 426 regime. *Ecology* 73: 2912-2207.

- 427 Brewin, P. A., S. T. Buckton, and S. J. Ormerod. 2000. The seasonal dynamics and
 428 persistence of stream macroinvertebrates in Nepal: do monsoon floods
 429 represent disturbance? *Freshwater Biology* 44: 581-594.
- 430 Brown, B.L. 2007. Habitat heterogeneity and disturbance influence patterns of
 431 community temporal variability in a small temperate stream. *Hydrobiologia*
 432 586: 93-106.
- 433 Bunn, S. E. and A. H. Arthington. 2002. Basic principles and ecological
 434 consequences of altered flow regimes for aquatic biodiversity. *Environmental*
 435 *Management* 30: 492-507.
- 436 Chantha, S., L. Cloutier, and A. Cattaneo. 2000. Epiphytic algae and invertebrates on
 437 aquatic mosses in a Quebec stream. *Archiv fur Hydrobiologie* 147: 143-160.
- 438 Cobb, D. G., T.D. Galloway, and J. F. Flannagan. 1992. Effects of discharge and
 439 substrate stability on density and species composition of stream insects.
 440 *Canadian Journal of Fisheries and Aquatic Sciences* 49: 1788-1795.
- 441 Collier, K. J. 2002. Effects of flow regulation and sediment flushing on instream
 442 habitat and benthic invertebrates in a New Zealand river influenced by a
 443 volcanic eruption. *River Research and Applications* 18: 213-226.

- 444 Death, R. G. 2007. The effects of floods on aquatic invertebrate communities. In
 445 Aquatic Insects: Challenges to Populations. Ed. J. Lancaster and R. A. Briers.
 446 Pgs. 103-121.
- 447 Death, R.G. 2010. Disturbance and riverine benthic communities: what has it
 448 contributed to general ecological theory? River Research and Applications
 449 26: 15-25.
- 450 Effenberger, M., J. Engel, S. Diehl, and C. D. Matthaei. 2008. Disturbance history
 451 influences the distribution of stream invertebrates by altering microhabitat
 452 parameters: a field experiment. *Freshwater Biology* 53: 996-1011.
- 453 Effenberger, M., G. Sailer, C.R. Townsend, and C.D. Matthaei. 2006. Local
 454 disturbance history and habitat parameters influence the microdistribution of
 455 stream invertebrates. *Freshwater Biology* 51: 312-332.
- 456 Fritz, K.M., and W.K. Dodds. 2004. Resistance and resilience of macroinvertebrate
 457 assemblages to drying in a tallgrass prairie stream. *Hydrobiologia* 527: 99-112.
- 458 Gray, L.J. 1981. Species composition and life histories of aquatic insects in a lowland
 459 Sonoran desert stream. *The American Midland Naturalist* 106: 229-242.
- 460 Grimm, N.B., and S.G. Fisher. 1989. Stability of periphyton and macroinvertebrates
 461 to disturbance by flash floods in a desert stream. *Journal of the North
 462 American Benthological Society* 8: 293-307.

- 463 Harrison, L.R., and E.A. Keller. 2007. Modeling forced pool-riffle hydraulics in a
 464 boulder-bed stream, southern California. *Geomorphology* 83: 232-248.
- 465 Hax, C.L., and S.W. Golladay. 1998. Flow disturbance of macroinvertebrates
 466 inhabiting sediments and woody debris in a prairie stream. *American Midland*
 467 *Naturalist* 1392: 210-223.
- 468 Haxton, T.J., and C.S. Findlay. 2008. Meta-analysis of the impacts of water
 469 management on aquatic communities. *Canadian Journal of Fisheries and*
 470 *Aquatic Sciences* 65: 437-447.
- 471 Hedges, L.V., J. Gurevitch, and P.S. Curtis. 1999. The meta-analysis of response
 472 ratios in experimental ecology. *Ecology* 80: 1150-1156.
- 473 Holomuzki, J.R., and B.J. Biggs. 2000. Taxon-specific responses to high-flow
 474 disturbance in streams: implications for population persistence. *Journal of the*
 475 *North American Benthological Society* 19: 670-679.
- 476 Imbert J.B., J.M. Gonzalez, A. Basaguren, and J. Pozo. 2005. Influence of inorganic
 477 substrata size, leaf litter and woody debris removal on benthic invertebrates
 478 resistance to floods in two contrasting headwater streams. *International*
 479 *Review of Hydrobiology* 90: 51-70.
- 480 Jowett, I. G. 1997. Instream flow methods: a comparison of approaches. *Regulated*
 481 *Rivers: Research & Management* 13: 115-127.

- 482 Kilbane, G.M., and J.R. Holomuzki. 2004. Spatial attributes, scale, and species traits
 483 determine caddisfly distributional responses to flooding. *Journal of the North*
 484 *American Benthological Society* 23: 480-493.
- 485 Konrad, C.P., J.D. Olden., D.A. Lytle, T. Melis, J.C. Schmidt, E. Bray, M. Freeman,
 486 K. Gido, N. Hemphill, M. Kennard., L. McMullen, M.C. Mims, M. Pyron, C.
 487 Robinson, and J.G. Williams. 2012. Large-scale flow experiments for
 488 managing rivers. *BioScience* 61: 948-959.
- 489 Lajeunesse, M.J. 2010. Achieving synthesis with meta-analysis by combining and
 490 comparing all available studies. *Ecology* 91: 2561-2564.
- 491 Lancaster, J. 1992. Diel variations in the effect of spates on mayflies
 492 (Ephemeroptera: Baetis). *Canadian Journal of Zoology* 70: 1696-1700.
- 493 Lapointe, M., B. Eaton, S. Driscoll, and C. Latulippe. 2000. Modeling the probability
 494 of salmonid egg pocket scour due to floods. *Canadian Journal of Fisheries and*
 495 *Aquatic Sciences* 57: 1120-1130.
- 496 Lytle D.A. 2000. Biotic and abiotic effects of flash flooding in a montane desert
 497 stream. *Archiv fur Hydrobiologie* 150: 85-100.
- 498 Lytle, D.A., J.D. Olden, and L.E. McMullen. 2008. Drought-escape behaviors of
 499 aquatic insects may be adaptations to highly variable flow regimes
 500 characteristic of desert rivers. *The Southwestern Naturalist* 53: 399-402.

- 501 Lytle, D.A., and N.J. White. 2007. Rainfall cues and flash-flood escape in desert
 502 stream insects. *Journal of Insect Behavior* 20: 413-423.
- 503 Maier, K.J. 2001. The influence of floods on benthic invertebrate populations in a
 504 Swiss mountain stream and their strategies of damage prevention. *Archiv fur*
 505 *Hydrobiologie* 150: 227-247.
- 506 Matthaei, C.D., C.J. Arbuckle, and C.R. Townsend. 2000. Stable surface stones as
 507 refugia for invertebrates during disturbance in a New Zealand stream. *Journal*
 508 *of the North American Benthological Society* 19: 82-93.
- 509 Matthaei, C.D., and H. Huber. 2002. Microform bed clusters: are they preferred
 510 habitats for invertebrates in a flood-prone stream? *Freshwater Biology*
 511 47:2174-2190.
- 512 Matthaei, C.D., U. Uehlinger, and A. Frutiger. 1997. Response of benthic
 513 invertebrates to natural versus experimental disturbance in a Swiss prealpine
 514 river. *Freshwater Biology* 37: 61-77.
- 515 Miller, A.M., and S.W. Golladay. 1996. Effects of spates and drying on
 516 macroinvertebrate assemblages of an intermittent and a perennial prairie
 517 stream. *Journal of the North American Benthological Society* 15: 670-689.

- 518 Monk, W.A., P.J. Wood, D.M. Hannah, D.A. Wilson, C.A. Extence, and R.P. Chadd.
 519 2006. Flow variability and macroinvertebrate community response within
 520 riverine systems. *River Research and Applications* 22: 595- 615.
- 521 Naiman, R. J., S. E. Bunn, C. Nilsson, G. E. Petts, G. Pinay, and L. C. Thompson.
 522 2002. Legitimizing fluvial ecosystems as users of water: an overview.
 523 *Environmental Management* 30: 455-467.
- 524 Negishi, J.N., M. Inou, and M. Nunokawa. 2002. Effects of channelization on stream
 525 habitat in relation to a spate and flow refugia for macroinvertebrates in
 526 northern Japan. *Freshwater Biology* 47: 1515-1529.
- 527 Negishi, J.N., and J. S. Richardson. 2006. An experimental test of the effects of food
 528 resources and hydraulic refuge on patch colonization by stream
 529 macroinvertebrates during spates. *Journal of Animal Ecology* 75: 118-129.
- 530 Nilsson, C., C. A. Reidy, M. Dynesius, and C. Revenga. 2005. Fragmentation and
 531 flow regulation of the world's large river systems. *Science* 308: 405-408.
- 532 Olsen, D.A., and C.R. Townsend. 2004. Flood effects on invertebrates, sediments
 533 and particulate organic matter in the hyporheic zone of a gravel-bed stream.
 534 *Freshwater Biology* 50: 839-853.

- 535 Orr, C.H., S.J. Kroiss, K.L. Rogers, and E.H. Stanley. 2008. Downstream benthic
 536 responses to small dam removal in a coldwater stream. *River Research and*
 537 *Applications* 24: 804-822.
- 538 Ortiz, J.D., and M.A. Puig. 2007. Point source effects on density, biomass, and
 539 diversity of benthic macroinvertebrates in a Mediterranean stream. *River*
 540 *Research and Applications* 23: 155-170.
- 541 Osenberg, C.W., O. Sarnelle, S.D. Cooper, and R.D. Holt. 1999. Resolving
 542 ecological questions through meta-analysis: goals, metrics and models.
 543 *Ecology* 80: 1105-1117.
- 544 Palmer, M.A., P. Arensburger, A.P. Martin, and D.W. Denman. 1996. Disturbance
 545 and patch-specific responses: the interactive effects of woody debris and floods
 546 on lotic invertebrates. *Oecologia* 105: 247-257.
- 547 Palmer, M.S., A.E. Bely, and K.E. Berg. 1992. Response of invertebrates to lotic
 548 disturbance: a test of the hyporheic refuge hypothesis. *Oecologia* 89: 182-194.
- 549 Palmer, A.R. 1999. Detecting publication bias in meta-analyses: a case of fluctuating
 550 asymmetry and sexual selection. *The American Naturalist* 154: 220-233.
- 551 Poff, N. L. 2009. Managing for variability to sustain freshwater ecosystems. *Journal*
 552 *of Water Resources Planning and Management* 135: 1-4.

- 553 Poff, N.L., and J.K.H. Zimmerman. 2010. Ecological responses to altered flow
 554 regimes: a literature review to inform the science and management of
 555 environmental flows. *Freshwater Biology* 55: 194-205.
- 556 Rader, R.B., N.J. Voelz, and J.V. Ward. 2008. Post-flood recovery of a
 557 macroinvertebrate community in a regulated river: resilience of an
 558 anthropogenically altered ecosystem. *Restoration Ecology* 16: 24-33.
- 559 Resh, V.H., A.V. Brown, A.P. Covich, M.E. Gurtz, H.W. Li, G.W. Minshall, S.R.
 560 Reice, A.L. Sheldon, J.B. Wallace, and R.C. Wissmar. 1988. The role of
 561 disturbance in stream ecology. *Journal of the North American Benthological*
 562 *Society* 7: 433-455.
- 563 Robinson, C.T., U. Uehlinger, and M.T. Monaghan. 2004. Stream ecosystem
 564 response to multiple experimental floods from a reservoir. *River Research and*
 565 *Applications* 20: 359-377.
- 566 Robinson, C.T., and U. Uehlinger. 2008. Experimental floods cause ecosystem
 567 regime shift in a regulated river. *Ecological applications* 18: 511-526.
- 568 Robson, B.J. 1996. Small spate disturbance and the complexity of habitat architecture
 569 in Mountain River, Tasmania. *Marine and Freshwater Research* 47: 851-855.

- 570 Rosenberg, M.S., D.C. Adams, and J. Gurevitch. 2000. MetaWin: Statistical
 571 Software for Meta-analysis. Version 2. Sinauer Associates, Sunderland,
 572 Massachusetts.
- 573 Scarsbrook, M.R., and C.D. Townsend. 1993. Stream community structure in relation
 574 to spatial and temporal variation: a habitat templet study of two contrasting
 575 New Zealand streams. *Freshwater Biology* 29: 395-4010.
- 576 Scrimgeour G.J., and M.J. Winterbourn. 1989. Effects of floods on epilithon and
 577 benthic macroinvertebrate populations in an unstable New Zealand river.
 578 *Hydrobiologia* 171: 33-44.
- 579 Shafroth, P.B., A.C. Wilcox, D.A. Lytle, J.T. Hickey, D.C. Andersen, V.B.
 580 Beauchamp, A. Hautzinger, L.E. McMullen, and A. Warner. 2010. Ecosystem
 581 effects of environmental flows: modeling and experimental floods in a dryland
 582 river. *Freshwater Biology* 55: 68-85.
- 583 Shurin, J.B., E.T. Borer, E.W. Seabloom, K. Anderson, C.A. Blanchette, B. Broitman,
 584 S.D. Cooper, and B.S. Halpern. 2002. A cross-ecosystem comparison of the
 585 strength of trophic cascades. *Ecology Letters* 5: 785-791.
- 586 SigmaPlot. 2004. Version 9.0. SYSTAT software.

- 587 Silver, P., D. Wooster, and M.A. Palmer. 2004. Chironomid responses to spatially
 588 structured, dynamic, streambed landscapes. *Journal of the North American*
 589 *Benthological Society* 23: 69-77.
- 590 Sokal, R.R. and F.J. Rohlf. 2000. *Biometry*. 3rd ed. W.H. Freeman and Company.
 591 Pgs. 251-252.
- 592 Souchon, Y., C. Sabaton, R. Deibel, D. Reiser, J. Kershner, M. Gard, C. Katopodis, P.
 593 Leonard, N.L. Poff, W.J. Miller, and B.L. Lamb. 2008. Detecting biological
 594 responses to flow management: missed opportunities, further directions. *River*
 595 *Research and Applications* 24: 506-518.
- 596 Stewart, G.B., H.R. Bayliss, D.A. Showler, W.J. Sutherland, and A.S. Pullin. 2009.
 597 Effectiveness of engineered instream structure mitigation measures to increase
 598 salmonid abundance: a systematic review. *Ecological Applications* 19: 931-
 599 941.
- 600 Stock, J.D., and I.J. Schlosser. 1991. Short-term effects of a catastrophic beaver dam
 601 collapse on a stream fish community. *Environmental Biology of Fishes* 31:
 602 123-129.
- 603 Tharme, R. E. 2003. A global perspective on environmental flow assessment:
 604 emerging trends in the development and application of environmental flow
 605 methodologies for rivers. *River Research and Applications* 19: 397-441.

606 Thiere, G., and R. Schulz. 2004. Runoff-related agricultural impact in relation to
 607 macroinvertebrate communities of the Lourens River, South Africa. *Water*
 608 *Research* 38: 3092-3102.

609 Thomson, J.R. 2002. The effects of hydrological disturbance on the densities of
 610 macroinvertebrate predators and their prey in a coastal stream. *Freshwater*
 611 *Biology* 47: 1333-1351.

612 Tummers, B. 2005-2006. Data Thief III. Software.

613 Wantzen, K.M. 1998. Effects of siltation on benthic communities in clear water
 614 streams in Mato Grosso, Brazil. *Verhandlungen Internationale Vereinigung*
 615 *für Theoretische und Angewandte Limnologie* 26: 1155-1159.

616 Yeston, J., R. Coontz, J. Smith, and C. Ash. 2006. A thirsty world. *Science* 313:
 617 1067.

618

619

620

621

622

623

624

625 **Table 1. Characteristics of all included studies.**

Reference	Country	River(s)	Invertebrates	Collection method	Flood type	Multi-sites?
Boulton et al. (1992)	United States (Arizona)	Sycamore Creek	Common taxa	Core	Natural	N
Bond and Downes (2000)	Australia	Steavenson	Hydropsychid caddisflies	Surber	Natural	N
Baumgartner and Waringer (1997)	Austria	Mauerbach	Overall abundance	Surber	Natural	Y
Angradi (1997)	United States (West Virginia)	Wilson Hollow Stream	Most abundant	Surber	Natural	N

Collier (2002)	Cobb et al. (1992)	Chantha et al. (2000)	Brown (2007)	Brewin et al. (2000)
New Zealand	Canada (Manitoba)	Canada (Quebec)	United States (New Hampshire)	Nepal
Tongagiro	Wilson Creek	Ruisseau Epinette	Alder Brook	Likhu Khola streams
<i>Deleatidium</i> and <i>Cricotopus</i>	Main groups	Abundance overall	Overall abundance	Together >90%
Surber	Hess	Hess	Metal frame	Surber
Managed	Natural	Natural	Natural	Natural
Y	Y	N	Y	N

Hax and Golladay (1998)	Fritz and Dodds (2004)	Effenberger et al. (2006)	Effenberger et al. (2008)
United States (Texas)	United States (Kansas)	New Zealand and Germany	Germany
Sister Grove Creek	Kings Creek tributaries	Kye Bum and Schmiedlaine	Eyach stream
Dominant taxa	>1mm total body length	Most common	5 most common
D-net	Stovepipe	Surber	Tile
Managed	Natural	Natural	Natural
Y	N	N	N

Lytle (2000)	Lancaster (1992)	Kilbane and Holomuzki (2004)	Imbert et al. (2005)	Holomuzki and Biggs (2000)
United States (Arizona)	Canada (BC)	United States (Ohio)	Spain	New Zealand
North Fork Cave Creek	Streamside channels at Mayfly Creek	Rocky Fork River tributary	Cuchillo and Salderrey streams	Laboratory flume
All	<i>Baetis</i>	2 numerically dominant caddisflies	10 predominant	Tested taxa
Box	Surber	Surber	Multiple	Visual
Natural	Managed	Natural	Natural	Managed
No	No	Yes	No	N

Negishi et al. (2002)	Miller and Golladay (1996)	Matthaei and Huber (2002)	Matthaei et al. (1997)	Matthaei et al. (2000)	Maier (2001)
Japan	United States (Oklahoma)	Germany	Switzerland	New Zealand	Switzerland
Nukanan Stream	Buncombe and Brier Creeks	Schmiedlaine	Necker River	Kye Burn	Kalte Sense
Overall abundance	Common	Common	Common	Common	5 dominant insects
Surber	Hess	Stones	Surber	Stones	Surber
Natural	Natural	Natural	Natural	Natural	Natural
Yes	No/Yes	No	No	No	No

Palmer et al. (1996)	United States (Virginia)	Goose Creek	Ortiz and Puig (2007)	Spain La Tordera	Orr et al. (2008)	United States (Wisconsin) Boulder Creek	Olsen and Townsend (2004)	New Zealand Kye Bum	Negishi and Richardson (2006)	Canada (BC) Spring Creek
Copepods and chironomids			Overall abundance		Major groups and trichopterans		Select taxa		Numerically dominant	
Core			Surber		Hess		Multiple		Cages	
Natural			Natural		Managed (Dam removal)		Natural		Natural	
No			No		No		No		No	

Robson (1996)	Robinson and Uehlinger (2008)	Robinson et al. (2004)	Rader et al. (2008)	Palmer et al. (1992)
Tasmania	Switzerland	Switzerland	United States (Colorado)	United States (Virginia)
Mountain River	Spol River	Spol River	Colorado River	Goose Creek
All	Common	Common	Overall abundance	Meiofauna
Quadrat	Hess	Hess	Surber	Core
Natural	Managed	Managed	Natural	Natural
Yes	No	Yes	No	No

Thiere and Schulz. (2004)	Stock and Schlosser (1991)	Silver et al. (2004)	Shafroth et al. (2010)	Scrimgeour and Winterbourn (1989)
South Africa	United States (Minnesota)	United States (Virginia)	United States (Arizona)	New Zealand
Lourens River	Gould Creek	Goose Creek	Bill Williams River	Ashley River
Common taxa	Insects overall	Chironomids	3 representative groups	Most common
Rocks	Surber	Leafpacks	D-net	Surber
Natural	Natural (Beaver dam)	Natural	Managed	Natural
No	Yes	Yes	Yes	No

Wantzen (1998)	Thomson (2002)
Brazil	Australia
Corrego	Cumberland River
Tenente	Common predators
Common	Electric pump
Artificial substrate	Natural
Natural	
Yes	No

preprint

Table 2. Characteristics of the two separate primary datasets used in meta-analyses.

	General dataset	Taxon-specific dataset
Sample unit	Before/ after flood abundance of total invertebrate count	Before/ after flood abundance of specific taxonomic units
Benefit	Minimize pseudoreplication within each study	All taxonomic groups from each study contribute equally to results
Bias to results	Taxa of highest abundance in each study have more influence	Higher in-study replication
Study inclusion criteria	Either: 1) Report total invertebrate abundance before/ after flood OR 2) Report abundance before/	Report abundance before/ after flood for at least one specific taxonomic group at any taxonomic level.

after flood for at least
three orders of
invertebrates (data will
be aggregated)

preprint

Table 3. P-values for categorical comparisons, sample sizes for all groups used in categorical comparisons, and results of Tukey-Kramer test for unplanned comparisons of group mean effect sizes for all categorical comparisons that exhibited significant differences among groups.

Group	General dataset			Taxon-specific dataset		
	<i>p</i>	<i>n</i>	T-K	<i>p</i>	<i>n</i>	T-K
Flood Type:	0.98			0.40		
Natural		78			242	
Managed		12			98	
Collection Method:	0.12			0.17		
Surber		20			68	
Hess		21			85	
Substrate		12			65	
Other		24			100	
Core		13			22	
Habitat:	<0.01			0.003		
Pool		5	a		24	c
Riffle		39	b		146	d
Run		8	b		30	e
Substrate:	0.63			0.003		
Gravel		32			105	f
Cobble		30			127	g
Boulder		n/a			28	f
Sand		8			31	f
Wood		n/a			14	g

Bedrock		9	29	f,g
Dry vs. Wet:	0.512		0.675	
Dry		30	134	
Wet		60	203	
Inverts- Ordinal or Higher:	n/a		0.26	
Coleoptera			20	
Eumalacostraca			15	
Annelida			22	
Ephemeroptera			70	
Diptera			76	
Trichoptera			49	
Plecoptera			46	
Acari			7	
Mollusca			8	
Platyhelminthes			9	
Ephemeroptera	n/a		0.72	
Baetidae			34	
Heptageniidae			21	
Leptophlebiidae			32	
Diptera:	n/a		0.049	
Ceratopogonidae			20	j,k
Chironomidae			83	k
Tipulidae			12	j
Simuliidae			24	j
Trichoptera:	n/a		0.705	
Hydropsychidae			11	
Lepidostomatidae			5	
Limnephilidae			10	

Plecoptera:

0.324

Nemouridae

20

Leuctridae

18

NOTES: *n* is the sample size. T-K stands for Tukey-Kramer. For the T-K results, groups with the same letter are not significantly different from each other.

esa
preprint

Figure 1. Effect size (natural log of invertebrate density post-floods/ invertebrate density pre-floods) of floods on aquatic invertebrate density and 95% confidence intervals. A scale for effect sizes as converted to percent reduction of invertebrates is on the right side of the figure. The black circles are effect sizes for sample units derived from the general data set, and the grey diamonds are effect sizes for sample units derived from the taxon-specific data set. The dashed line at 0 indicates which effect size results are significant; those with confidence intervals overlapping the dotted line are not significant. The overall (cumulative) effect size is shown, as well as effect sizes estimated from categorical analyses of flood type, collection method, habitat type, substrate type, and whether the flood happened in a 'wet' or 'dry' month.

Figure 2. Effect size (natural log of invertebrate density post-floods/ invertebrate density pre-floods) of floods on overall aquatic invertebrate density versus time since the flood event, within the first 10 days of a flood event.

Figure 3. Effect size (natural log of invertebrate density post-floods/ invertebrate density pre-floods) of floods on overall aquatic invertebrate density versus relative flood magnitude, for riffle or run habitats composed of primarily cobble or gravel substrate.

Figure 4. Effect size (natural log of invertebrate density post-floods/ invertebrate density pre-floods) of floods on aquatic invertebrate density of different taxonomic groups and 95% confidence intervals. A scale for effect sizes as converted to percent reduction of invertebrates is on the right side of the figure. The dashed line at 0 indicates which effect size results are significant (the effect of floods on density of these groups was significant); those that have confidence intervals overlapping the dotted line were not significant. Results from categorical

analyses that were conducted at lower taxonomic levels are boxed along with the effect size estimated for their parent group.

esa

preprint




Figure 1.




Figure 2.




Figure 3.




Figure 4.